

## Subject Index

### Volume 62 (1992)

- <sup>13</sup>C, high resolution NMR spectra, fish lipids, *n*-3, *n*-6, *n*-7, *n*-9, eicosapentaenoic acid, docosahexaenoic acid, arachidonic acid, 239
- aggregation properties, gangliosides, ceramide, lipid moiety, critical micellar concentration, granule cells, 1
- aging of liposomes, leak-in rate, permeability, physical stability, fluid-state bilayers, lyso-phosphatidylcholine effect, cholesterol effect, 113
- air-water interface, fluorescent lipid probes, monolayers, phosphatidylcholine, phosphatidylglycerol, gangliosides, 293
- allylic dihydroxy unsaturated fatty ester, dehydration, 177
- aminoglycoside, deuterium, interaction, nuclear magnetic resonance, phosphatidylcholine, phosphatidylinositol, 153
- antioxidants, fluorescence, bilayers, quenching, energy transfer, 123
- apoprotein, cholesterol, interaction, guanidine group, arginine, hydroxyl group, 229
- arachidonic, linolenic, radical, singlet oxygen, photo-oxidation, high performance liquid chromatography (HPLC), 165
- arginine, apoprotein, cholesterol, interaction, guanidine group, hydroxyl group, 229
- bilayer phase transition, DHPC, vesicle aggregation,  $\text{Ca}^{2+}$ , ethanol, stereoisomers, 253
- bilayers, antioxidants, fluorescence, quenching, energy transfer, 123
- 1,2-bis-(tricosano 10,12 diynoyl)-*rac*-glycero-3-(*p*-toluene-sulfonate), polymerizable lipid, Raman scattering, resonance Raman, diacetylene, crystallinity, 139
- BNCT, ether lipid, boron neutron capture therapy, phosphoamidite, glycidyl ether, 185
- boron neutron capture therapy, ether lipid, BNCT, phosphoamidite, glycidyl ether, 185
- $\text{Ca}^{2+}$ , DHPC, vesicle aggregation, bilayer phase transition, ethanol, stereoisomers, 253
- calcium antagonists, membrane perturbation, EPR spectroscopy, 11
- ceramide, conformation, 2,3-dihydroxy fatty acid, phytosphingosine, hydrogen bonds, 79
- ceramide, gangliosides, lipid moiety, aggregation properties, critical micellar concentration, granule cells, 1
- changes in membrane fluidity, MPTP (1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine), MPDP<sup>+</sup> (1-methyl-4-phenyl-2,3-dihydropyridinium), MPP<sup>+</sup> (1-methyl-4-phenylpyridinium), cytochrome P-450 inhibition, 303
- cholera toxin, resonance energy transfer, glycosphingolipid clustering, ganglioside GM1, 4-nitrobenz-2-oxa-1,3,-diazole (NBD), diphenylhexatriene (DPH), 269
- cholesterol effect, leak-in rate, permeability, physical stability, fluid-state bilayers, aging of liposomes, lyso-phosphatidylcholine effect, 113
- cholesterol hydroperoxides, oxysterols, cholesterol oxidation, oxidized LDL, 209
- cholesterol oxidation, oxysterols, cholesterol hydroperoxides, oxidized LDL, 209
- cholesterol, apoprotein, interaction, guanidine group, arginine, hydroxyl group, 229
- conformation, ceramide, 2,3-dihydroxy fatty acid, phytosphingosine, hydrogen bonds, 79
- conformational analysis, 15-oxygenated sterols, inhibitors of sterol synthesis, <sup>1</sup>H- and <sup>13</sup>C-NMR of sterols, 55
- counterion-headgroup interaction, diacetylenic phospholipids, tubules, morphology modulation, surface-charge screening, monolayer films, 193
- critical micellar concentration, gangliosides, ceramide, lipid moiety, aggregation properties, granule cells, 1
- crystallinity, polymerizable lipid, Raman scattering, resonance Raman, diacetylene, 1,2-bis-(tricosano 10,12 diynoyl)-*rac*-glycero-3-(*p*-toluene-sulfonate), 31
- cysteamine, peroxidation, liposomes, phospholipids, 31
- cytochrome P-450 inhibition, MPTP (1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine), MPDP<sup>+</sup> (1-methyl-4-phenyl-2,3-dihydropyridinium), MPP<sup>+</sup> (1-methyl-4-phenylpyridinium), changes in membrane fluidity, 303
- dehydration, allylic dihydroxy unsaturated fatty ester, 177
- deuterium, aminoglycoside, interaction, nuclear magnetic resonance, phosphatidylcholine, phosphatidylinositol, 153
- DHPC, vesicle aggregation, bilayer phase transition,  $\text{Ca}^{2+}$ , ethanol, stereoisomers, 253
- diacetylene, polymerizable lipid, Raman scattering, resonance Raman, crystallinity, 1,2-bis-(tricosano 10,12 diynoyl)-*rac*-glycero-3-(*p*-toluene-sulfonate), 139
- diacetylenic phospholipids, tubules, counterion-headgroup interaction, morphology modulation, surface-charge screening, monolayer films, 193
- diacylglycerol, phospholipid vesicles, fusion, infrared spectroscopy, <sup>31</sup>P-nuclear magnetic resonance, 215
- 2,3-dihydroxy fatty acid, ceramide, conformation, phytosphingosine, hydrogen bonds, 79
- diphenylhexatriene (DPH), resonance energy transfer, glycosphingolipid clustering, cholera toxin, ganglioside GM1, 4-nitrobenz-2-oxa-1,3,-diazole (NBD), 269

- 2**
- DNA gyrase, two-dimensional crystallization, protein, phospholipid, novobiocin, synthesis, 93  
 docosahexaenoic acid, high resolution NMR spectra,  $^{13}\text{C}$ , fish lipids, n-3, n-6, n-7, n-9, eicosapentaenoic acid, arachidonic acid, 239  
 domain, methylation, phospholipids, monolayer, fluorescence microscopy, phase transition, 69  
 dynamic light scattering, liposome characterization, small angle X-ray scattering, encapsulated volume, size distribution, 105  
 eicosapentaenoic acid, high resolution NMR spectra,  $^{13}\text{C}$ , fish lipids, n-3, n-6, n-7, n-9, docosahexaenoic acid, arachidonic acid, 239  
 encapsulated volume, liposome characterization, dynamic light scattering, small angle X-ray scattering, size distribution, 105  
 energy transfer, antioxidants, fluorescence, bilayers, quenching, 123  
 EPR spectroscopy, calcium antagonists, membrane perturbation, 11  
 ethanol, DHPC, vesicle aggregation, bilayer phase transition,  $\text{Ca}^{2+}$ , stereoisomers, 253  
 ether lipid, boron neutron capture therapy, BNCT, phosphoamidite, glycidyl ether, 185  
 excimer, pyrene-labeled lipid, phase transition, 39  
 fatty ester, thiolsulfinate, synthesis, physical properties, 145  
 fish lipids, high resolution NMR spectra,  $^{13}\text{C}$ , n-3, n-6, n-7, n-9, eicosapentaenoic acid, docosahexaenoic acid, arachidonic acid, 239  
 fluid-state bilayers, leak-in rate, permeability, physical stability, aging of liposomes, lyso-phosphatidylcholine effect, cholesterol effect, 113  
 fluorescence microscopy, methylation, phospholipids, monolayer, domain, phase transition, 69  
 fluorescence, antioxidants, bilayers, quenching, energy transfer, 123  
 fluorescent lipid probes, air-water interface, monolayers, phosphatidylcholine, phosphatidylglycerol, gangliosides, 293  
 Fourier transform infrared spectroscopy, palmitic acid, phosphatidylcholine, isotopic substitution, 19  
 fusion, diacylglycerol, phospholipid vesicles, infrared spectroscopy,  $^{31}\text{P}$ -nuclear magnetic resonance, 215  
 ganglioside GM1, resonance energy transfer, glycosphingolipid clustering, cholera toxin, 4-nitrobenz-2-oxa-1,3-diazole (NBD), diphenylhexatriene (DPH), 269  
 gangliosides, ceramide, lipid moiety, aggregation properties, critical micellar concentration, granule cells, 1  
 gangliosides, fluorescent lipid probes, air-water interface, monolayers, phosphatidylcholine, phosphatidylglycerol, 293  
 glycidyl ether, ether lipid, boron neutron capture therapy, BNCT, phosphoamidite, 185  
 glycolipid, mannose mycolate, *Rhodococcus*, mycolic acid, 311  
 glycosphingolipid clustering, resonance energy transfer, cholera toxin, ganglioside GM1, 4-nitrobenz-2-oxa-1,3-diazole (NBD), diphenylhexatriene (DPH), 269  
 granule cells, gangliosides, ceramide, lipid moiety, aggregation properties, critical micellar concentration, 1  
 growth rate, myelin figure, phosphatidylcholine, phosphatidylethanolamine, 87  
 guanidine group, apoprotein, cholesterol, interaction, arginine, hydroxyl group, 229  
 $^1\text{H}$ - and  $^{13}\text{C}$ -NMR of sterols, 15-oxygenated sterols, inhibitors of sterol synthesis, conformational analysis, 55  
 $^1\text{H}$ -NMR, local anesthetics, lipid bilayer, nuclear Overhauser effect, molecular dynamics, 45  
 high performance liquid chromatography (HPLC), linolenic, arachidonic, radical, singlet oxygen, photo-oxidation, 165  
 high resolution NMR spectra,  $^{13}\text{C}$ , fish lipids, n-3, n-6, n-7, n-9, eicosapentaenoic acid, docosahexaenoic acid, arachidonic acid, 239  
 hydrogen bonds, ceramide, conformation, 2,3-dihydroxy fatty acid, phytosphingosine, 79  
 hydroxyl group, apoprotein, cholesterol, interaction, guanidine group, arginine, 229  
 infrared spectroscopy, diacylglycerol, phospholipid vesicles, fusion,  $^{31}\text{P}$ -nuclear magnetic resonance, 215  
 inhibitors of sterol synthesis, 15-oxygenated sterols,  $^1\text{H}$ - and  $^{13}\text{C}$ -NMR of sterols, conformational analysis, 55  
 interaction, aminoglycoside, deuterium, nuclear magnetic resonance, phosphatidylcholine, phosphatidylinositol, 153  
 interaction, apoprotein, cholesterol, guanidine group, arginine, hydroxyl group, 229  
 isotopic substitution, palmitic acid, phosphatidylcholine, Fourier transform infrared spectroscopy, 19  
 leak-in rate, permeability, physical stability, fluid-state bilayers, aging of liposomes, lyso-phosphatidylcholine effect, cholesterol effect, 113  
 linolenic, arachidonic, radical, singlet oxygen, photo-oxidation, high performance liquid chromatography (HPLC), 165  
 lipid bilayer, local anesthetics, nuclear Overhauser effect,  $^1\text{H}$ -NMR, molecular dynamics, 45  
 lipid moiety, gangliosides, ceramide, aggregation properties, critical micellar concentration, granule cells, 1  
 liposome characterization, dynamic light scattering, small angle X-ray scattering, encapsulated volume, size distribution, 105  
 liposomes, peroxidation, phospholipids, cysteamine, 31  
 local anesthetics, lipid bilayer, nuclear Overhauser effect,  $^1\text{H}$ -NMR, molecular dynamics, 45  
 lyso-phosphatidylcholine effect, leak-in rate, permeability, physical stability, fluid-state bilayers, aging of liposomes, cholesterol effect, 113  
 mannosyl mycolate, *Rhodococcus*, mycolic acid, glycolipid, 311  
 membrane perturbation, calcium antagonists, EPR spectroscopy, 11

- methylation, phospholipids, monolayer, domain, fluorescence microscopy, phase transition, 69
- microwave, transesterification, pyrrolidides, 281
- molecular dynamics, local anesthetics, lipid bilayer, nuclear Overhauser effect,  $^1\text{H-NMR}$ , 45
- monolayer films, diacylenic phospholipids, tubules, counterion-headgroup interaction, morphology modulation, surface-charge screening, 193
- monolayer, methylation, phospholipids, domain, fluorescence microscopy, phase transition, 69
- monolayers, fluorescent lipid probes, air-water interface, phosphatidylcholine, phosphatidylglycerol, gangliosides, 293
- morphology modulation, diacylenic phospholipids, tubules, counterion-headgroup interaction, surface-charge screening, monolayer films, 193
- $\text{MPDP}^+$  (1-methyl-4-phenyl-2,3-dihydropyridinium), MPTP (1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine),  $\text{MPP}^+$  (1-methyl-4-phenylpyridinium), changes in membrane fluidity, cytochrome  $P-450$  inhibition, 303
- $\text{MPP}^+$  (1-methyl-4-phenylpyridinium), MPTP (1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine),  $\text{MPDP}^+$  (1-methyl-4-phenyl-2,3-dihydropyridinium), changes in membrane fluidity, cytochrome  $P-450$  inhibition, 303
- MPTP (1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine),  $\text{MPDP}^+$  (1-methyl-4-phenyl-2,3-dihydropyridinium),  $\text{MPP}^+$  (1-methyl-4-phenylpyridinium), changes in membrane fluidity, cytochrome  $P-450$  inhibition, 303
- mycolic acid, mannose mycolate, *Rhodococcus*, glycolipid, 311
- myelin figure, growth rate, phosphatidylcholine, phosphatidylethanolamine, 87
- n*-3, high resolution NMR spectra,  $^{13}\text{C}$ , fish lipids, *n*-6, *n*-7, *n*-9, eicosapentaenoic acid, docosahexaenoic acid, arachidonic acid, 239
- n*-6, high resolution NMR spectra,  $^{13}\text{C}$ , fish lipids, *n*-3, *n*-7, *n*-9, eicosapentaenoic acid, docosahexaenoic acid, arachidonic acid, 239
- n*-7, high resolution NMR spectra,  $^{13}\text{C}$ , fish lipids, *n*-3, *n*-6, *n*-9, eicosapentaenoic acid, docosahexaenoic acid, arachidonic acid, 239
- n*-9, high resolution NMR spectra,  $^{13}\text{C}$ , fish lipids, *n*-3, *n*-6, *n*-7, eicosapentaenoic acid, docosahexaenoic acid, arachidonic acid, 239
- 4-nitrobenz-2-oxa-1,3-diazole (NBD), resonance energy transfer, glycosphingolipid clustering, cholera toxin, ganglioside GM1, diphenylhexatriene (DPH), 269
- novobiocin, two-dimensional crystallization, protein, phospholipid, synthesis, DNA gyrase, 93
- nuclear magnetic resonance, aminoglycoside, deuterium, interaction, phosphatidylcholine, phosphatidylinositol, 153
- nuclear Overhauser effect, local anesthetics, lipid bilayer,  $^1\text{H-NMR}$ , molecular dynamics, 45
- oxidized LDL, oxysterols, cholesterol hydroperoxides, cholesterol oxidation, 209
- 15-oxygenated sterols, inhibitors of sterol synthesis,  $^1\text{H}$ - and  $^{13}\text{C-NMR}$  of sterols, conformational analysis, 55
- oxysterols, cholesterol hydroperoxides, cholesterol oxidation, oxidized LDL, 209
- $^{31}\text{P}$ -nuclear magnetic resonance, diacylglycerol, phospholipid vesicles, fusion, infrared spectroscopy, 215
- palmitic acid, phosphatidylcholine, Fourier transform infrared spectroscopy, isotopic substitution, 19
- permeability, leak-in rate, physical stability, fluid-state bilayers, aging of liposomes, lyso-phosphatidylcholine effect, cholesterol effect, 113
- peroxidation, liposomes, phospholipids, cysteamine, 31
- phase transition, methylation, phospholipids, monolayer, domain, fluorescence microscopy, 69
- phase transition, pyrene-labeled lipid, excimer, 39
- phenosafranin dye, phospholipid liposome, thionine dye, photovoltage, photoelectrochemistry of dye, 225
- phosphatidylcholine, aminoglycoside, deuterium, interaction, nuclear magnetic resonance, phosphatidylinositol, 153
- phosphatidylcholine, fluorescent lipid probes, air-water interface, monolayers, phosphatidylglycerol, gangliosides, 293
- phosphatidylcholine, myelin figure, growth rate, phosphatidylethanolamine, 87
- phosphatidylcholine, palmitic acid, Fourier transform infrared spectroscopy, isotopic substitution, 19
- phosphatidylethanolamine, myelin figure, growth rate, phosphatidylcholine, 87
- phosphatidylglycerol, fluorescent lipid probes, air-water interface, monolayers, phosphatidylcholine, gangliosides, 293
- phosphatidylinositol, aminoglycoside, deuterium, interaction, nuclear magnetic resonance, phosphatidylcholine, 153
- phosphoamidite, ether lipid, boron neutron capture therapy, BNCT, glycidyl ether, 185
- phospholipase C, phosphonate, phospholipase D, 263
- phospholipase D, phosphonate, phospholipase C, 263
- phospholipid liposome, phenosafranin dye, thionine dye, photovoltage, photoelectrochemistry of dye, 225
- phospholipid vesicles, diacylglycerol, fusion, infrared spectroscopy,  $^{31}\text{P}$ -nuclear magnetic resonance, 215
- phospholipid, two-dimensional crystallization, protein, novobiocin, synthesis, DNA gyrase, 93
- phospholipids, methylation, monolayer, domain, fluorescence microscopy, phase transition, 69
- phospholipids, peroxidation, liposomes, cysteamine, 31
- phosphonate, phospholipase C, phospholipase D, 263
- photo-oxidation, linolenic, arachidonic, radical, singlet oxygen, high performance liquid chromatography (HPLC), 165
- photoelectrochemistry of dye, phospholipid liposome, phenosafranin dye, thionine dye, photovoltage, 225
- photovoltage, phospholipid liposome, phenosafranin dye, thionine dye, photoelectrochemistry of dye, 225
- physical properties, thiolsulfinate, fatty ester, synthesis, 145
- physical stability, leak-in rate, permeability, fluid-state bilayers, aging of liposomes, lyso-phosphatidylcholine effect, cholesterol effect, 113
- phytosphingosine, ceramide, conformation, 2,3-dihydroxy fatty acid, hydrogen bonds, 79
- polymerizable lipid, Raman scattering, resonance Raman,

- diacetylene, crystallinity, 1,2-bis-(tricos 10,12 diynoyl)-*rac*-glycero-3-(*p*-toluene-sulfonate), 139
- protein, two-dimensional crystallization, phospholipid, novobiocin, synthesis, DNA gyrase, 93
- pyrene-labeled lipid, excimer, phase transition, 39
- pyrrolidides, microwave, transesterification, 281
- quenching, antioxidants, fluorescence, bilayers, energy transfer, 123
- radical, linolenic, arachidonic, singlet oxygen, photo-oxidation, high performance liquid chromatography (HPLC), 165
- Raman scattering, polymerizable lipid, resonance Raman, diacetylene, crystallinity, 1,2-bis-(tricos 10,12 diynoyl)-*rac*-glycero-3-(*p*-toluene-sulfonate), 139
- resonance energy transfer, glycosphingolipid clustering, cholera toxin, ganglioside GM1, 4-nitrobenz-2-oxa-1,3-diazole (NBD), diphenylhexatriene (DPH), 269
- resonance Raman, polymerizable lipid, Raman scattering, diacetylene, crystallinity, 1,2-bis-(tricos 10,12 diynoyl)-*rac*-glycero-3-(*p*-toluene-sulfonate), 139
- Rhodococcus*, mannose mycolate, mycolic acid, glycolipid, 311
- singlet oxygen, linolenic, arachidonic, radical, photo-oxidation, high performance liquid chromatography (HPLC), 165
- size distribution, liposome characterization, dynamic light scattering, small angle X-ray scattering, encapsulated volume, 105
- small angle X-ray scattering, liposome characterization, dynamic light scattering, encapsulated volume, size distribution, 105
- stereoisomers, DHPC, vesicle aggregation, bilayer phase transition,  $\text{Ca}^{2+}$ , ethanol, 253
- surface-charge screening, diacetylenic phospholipids, tubules, counterion-headgroup interaction, morphology modulation, monolayer films, 193
- synthesis, thiolsulfinate, fatty ester, physical properties, 145
- synthesis, two-dimensional crystallization, protein, phospholipid, novobiocin, DNA gyrase, 93
- thiolsulfinate, fatty ester, synthesis, physical properties, 145
- thionine dye, phospholipid liposome, phenosafranin dye, photovoltage, photoelectrochemistry of dye, 225
- transesterification, microwave, pyrrolidides, 281
- tubules, diacetylenic phospholipids, counterion-headgroup interaction, morphology modulation, surface-charge screening, monolayer films, 193
- two-dimensional crystallization, protein, phospholipid, novobiocin, synthesis, DNA gyrase, 93
- vesicle aggregation, DHPC, bilayer phase transition,  $\text{Ca}^{2+}$ , ethanol, stereoisomers, 253

## Author Index

### Volume 62 (1992)

Allevi, P.	209	Jarrell, H.C.	153
Antes, P.	269	Kézdy, F.J.	123
Aranda, F.J.	215	Kim, H-S.	55
Aursand, M.	239	Kinnunen, P.K.J.	253
Bakare, O.	145	Kirkpatrick, N.D.	55
Banerjee, P.	281	Klimov, A.N.	229
Bassi, R.	1	Klueva, N.N.	229
Behroozi, F.	139	Köiv, A.	253
Belova, E.V.	229	Kozhevnikova, K.A.	229
Bergelson, L.D.	293	Kuroda, Y.	45
Bhowmik, B.B.	225		
Bouwstra, J.	105	Lebeau, L.	93
Bredlow, A.	293	Lemmen, P.	185
Butko, P.	39	Lie Ken Jie, M.S.F.	145, 177
Carrier, D.	153	Malavasi, B.	209
Catapano, A.L.	209	Malik, S.	281
Cervato, G.	31	Markowitz, M.A.	193
Cestaro, B.	31	McKenna, R.L.	123
Chancón, J.N.	165	McLearie, J.	165
Cheng, K.H.	39	Micol, V.	215
Crommelin, D.J.A.	105, 113	Mingeot-Leclercq, M.-P.	153
Daniel, L.W.	263	Mioskowski, C.	93
Das, N.P.	303	Mishima, K.	87
Dasgupta, A.	281	Moochhala, S.M.	303
Emilsson, A.	263	Nakagawa, T.	45
Epps, D.E.	123	Nixon, A.B.	263
Fogliatto, R.	209	Ogihara, T.	87
Fujiwara, Y.	45	Olland, S.	93
Gaber, B.P.	139	Ondriaš, K.	11
Galla, H.J.	293	Ondriášová, E.	11
Gallii, G.	209	Ortiz, A.	215
Gatti, P.	31	Oudemans, P.	93
Gómez-Fernández, J.C.	19, 215	Pascher, I.	79
Gooris, G.	105	Pierson, T.S.	123
Grasdalen, H.	239	Pinkerton, F.D.	55
Grit, M.	113	Rasetti, M.F.	209
Han, F.	123	Roma, P.	209
Hinzmann, J.S.	123	Salomons, M.A.	105
Hui, S-W.	69	San Martin, C.	215
Hybl, A.	139	Sandhoff, K.	269
Hylarides, M.D.	55	Sato, T.	303
Ioneda, T.	311	Satoh, K.	87
		Schanck, A.	153

Schmitt, J.D.	263	Talsma, H.	105
Schnur, J.M.	193	Tomita, M.	87
Schoen, P.E.	139	Tulkens, P.M.	153
Schroepfer, Jr. G.J.	55		
Schwarzmann, G.	269		
Sen Verma, C.	225	Viani, P.	31
Shahi, G.S.	303	Villalain, J.	19
Sinclair, R.S.	165	Villalain, J.	215
Singh, A.	139, 193		
Smith, F.J.	165	Wakita, M.	45
Smith, I.C.P.	153	Werner, B.	185
Sonnino, S.	1	Wilson, W.K.	55
Staško, A.	11	Wong, K.P.	177
Steenbergen, M.J.	105	Wykle, R.L.	263
Sunamoto, J.	303		
Sundell, S.	79		
Swaminathan, S.	55	Yu, H.	69

